

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
<p>The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to the Department of Defense, Executive Service Directorate (0704-0188). Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.</p> <p><b>PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ORGANIZATION.</b></p>					
1. REPORT DATE (DD-MM-YYYY) 21/03/2017		2. REPORT TYPE final		3. DATES COVERED (From - To) 11/02/2013-21/03/2017	
4. TITLE AND SUBTITLE Rapid Extremity Pain Relief by Battlefield Acupuncture after Orthopedic Surgery: A Randomized Clinical Trial.				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Paul Crawford, MD				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Clinical Investigation Program Mike O'Callaghan Federal Medical Center 4700 Las Vegas Blvd North Nellis AFB, NV 89191				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Clinical Investigation Program Mike O'Callaghan Federal Medical Center 4700 Las Vegas Blvd North Nellis AFB, NV 89191				10. SPONSOR/MONITOR'S ACRONYM(S)  FDG201234H	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S) FDG201234H	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution is unlimited.					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT See attached					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT	b. ABSTRACT	c. THIS PAGE			Jill Clark
U	U	U	UU		19b. TELEPHONE NUMBER (Include area code) (702) 653-3298



**DEPARTMENT OF THE AIR FORCE AND  
DEPARTMENT OF VETERANS AFFAIRS**  
MIKE O'CALLAGHAN FEDERAL MEDICAL CENTER  
NELLIS AIR FORCE BASE, NEVADA



**FINAL REPORT**

**Project Title:**

Rapid Extremity Pain Relief by Battlefield Acupuncture after Orthopedic Surgery: A Randomized Clinical  
Trial.

**Project#:**

FDG201234H

**Date Submitted:**

21 March 2017

**Reporting Period:**

11 February 2013-21 March 2017

**Prepared for:**

Defense Technical Information Center  
ATTN: DTIC-OA  
8725 John J. Kingman Rd  
Fort Belvoir, VA 22060-6218

**Submitted by:**

Jill M. Clark, MBA/HCM, CCRP, CCRC  
Senior Research Associate/Research Manager  
Clinical Investigation Program  
Mike O'Callaghan Federal Medical Center (MOFMC)  
4700 Las Vegas Blvd North, Bldg 1300, Room 1884  
Nellis AFB, NV 89191  
Phone: (702) 653-3298  
Email: Jill.Clark.3.ctr@us.af.mil

**Distribution Statement A:**

Approved for public release: distribution unlimited.

**UNCLASSIFIED**

**Background:** Due to the emergence of improvised explosive devices (IED's) and the widespread use of body armor, there have been a higher proportion of orthopedic injuries from secondary blast injury than in previous conflicts. Since 2003, it is estimated that approximately 32,195 soldiers have been wounded in combat in the Iraq conflict alone. According to the American Academy of Orthopedic Surgeons, more than four and one half million knee arthroscopies and total knee replacements are performed worldwide each year in addition to millions of foot and ankle surgeries. These invasive procedures result in swelling and pain. The side effects of the pain medications are well known and a decrease in their use could prevent adverse effects of sedation and decreased job performance. Auricular acupuncture has been evaluated in multiple trials, and although generally proven to be useful, these trials were not rigorous. This study seeks to determine if modified Battlefield Acupuncture is more effective at relieving acute extremity pain, reducing medication use, decreasing time to full ambulation and improving quality of life than placebo acupuncture or standard care after lower extremity surgery.

**Methods:** We conducted a multi-site 3-arm randomized, double blind controlled trial of standard care alone versus standard care + placebo auricular acupuncture with ASP needles versus standard care + battlefield acupuncture with semi-permanent needles. We recruited subjects at the pre-operative visit within 2 Air Force orthopedics clinics for lower extremity surgery (knee or below). After consent, we randomized them to the above interventions. After surgery, within 30 minutes of arrival in the post-operative care unit, subjects received their designated intervention. Acupuncture was performed by physician acupuncturists. Subjects reported pain level immediately after acupuncture, 24, 28 168 and 720 hours later to a blinded research associate. Additionally, subjects completed a PIQ-6 30 days post-operatively, and opioid use was tracked for 30 days post-operatively. Statistical analysis with ANOVA, Pearson's correlation, Chi-square and Fisher's exact test, and multivariate analyses were performed on the data.

**Results:** We enrolled 233 Department of Defense beneficiaries >18 years old (92 females and 141 males) with a mean age of 44.5 years at 2 Air Force Medical Centers. We randomized 81 to modified BFA, 74 to placebo acupuncture and 78 to standard care. Overall pain levels were unchanged at each time point between groups. Subjects with worse pain were noted to take more opioid medication, and subjects older than 50 years who received modified BFA took 398 fewer morphine equivalent units than those who received standard therapy ( $p=.09$ )

**Conclusion:** The use of modified battlefield acupuncture protocol does not change pain over 30 days, but both modified BFA and sham acupuncture may reduce use of opioid analgesics in those >50 years of age.